



TITLE:

Fractional integrals and their commutators on martingale Orlicz spaces (The generalization of function spaces and its environment)

AUTHOR(S):

Arai, Ryutaro; Nakai, Eiichi; Sadasue, Gaku

CITATION:

Arai, Ryutaro ...[et al]. Fractional integrals and their commutators on martingale Orlicz spaces (The generalization of function spaces and its environment). 数理解析研究所講究録 2019, 2143: 12-22

ISSUE DATE:

2019-12

URL:

<http://hdl.handle.net/2433/254965>

RIGHT:

Fractional integrals and their commutators on martingale Orlicz spaces

茨城大学大学院理工学研究科 新井 龍太郎

Ryutaro Arai

Department of Mathematics, Ibaraki University

茨城大学大学院理工学研究科 中井 英一

Eiichi Nakai

Department of Mathematics, Ibaraki University

大阪教育大学教育学部 貞末 岳

Gaku Sadasue

Department of Mathematics, Osaka Kyoiku University

1 Introduction

This is an announcement of [2].

It is well known as the Hardy-Littlewood-Sobolev theorem that the fractional integral operators I_α on the Euclidean space \mathbb{R}^n is bounded from L_p to L_q for $1 < p < q < \infty$, $0 < \alpha < n$ and $-n/p + \alpha = -n/q$. For any BMO function b , Chanillo [4] proved the same boundedness of the commutator $[b, I_\alpha]$. Paluszyński [19] proved that, for any β -Lipschitz function b , $0 < \beta < 1$, the commutator $[b, I_\alpha]$ is bounded from L_p to L_q for $-n/p + \alpha + \beta = -n/q$ and from L_p to the Triebel-Lizorkin space $\dot{F}_{p,\infty}^\beta$.

In martingale theory, based on the result by Watari [23, Theorem 1.1], Chao and Ombe [5] proved the boundedness of the fractional integrals for H_p , L_p , BMO and Lipschitz spaces of the dyadic martingales. These fractional integrals were defined for more general martingales in [14, 20] and studied in [6, 15, 16]. In this paper we investigate the fractional integrals on martingale Orlicz spaces.

Let (Ω, \mathcal{F}, P) be a probability space and let $\{\mathcal{F}_n\}_{n \geq 0}$ be a nondecreasing sequence of sub- σ -algebras of \mathcal{F} such that $\mathcal{F} = \sigma(\bigcup_n \mathcal{F}_n)$. We suppose that every σ -algebra \mathcal{F}_n is generated by countable atoms, where $B \in \mathcal{F}_n$ is called an atom (more precisely a (\mathcal{F}_n, P) -atom), if any $A \subset B$ with $A \in \mathcal{F}_n$ satisfies $P(A) = P(B)$ or $P(A) = 0$. Denote by $A(\mathcal{F}_n)$ the set of all atoms in \mathcal{F}_n . The

expectation operator and the conditional expectation operators relative to \mathcal{F}_n are denoted by E and E_n , respectively.

We say a sequence $(f_n)_{n \geq 0}$ in L_1 is a martingale relative to $\{\mathcal{F}_n\}_{n \geq 0}$ if it is adapted to $\{\mathcal{F}_n\}_{n \geq 0}$ and satisfies $E_n[f_m] = f_n$ for every $n \leq m$. It is known as the Doob theorem that, if $p \in (1, \infty)$, then any L_p -bounded martingale converges in L_p . Moreover, if $p \in [1, \infty)$, then, for any $f \in L_p$, its corresponding martingale $(f_n)_{n \geq 0}$ with $f_n = E_n f$ is an L_p -bounded martingale and converges to f in L_p (see for example [17]). For this reason a function $f \in L_1$ and the corresponding martingale $(f_n)_{n \geq 0}$ will be denoted by the same symbol f .

We first recall the definition of generalized fractional integrals of martingales.

Definition 1.1 ([16]). Let $(\gamma_n)_{n \geq 0}$ be a non-increasing sequence of non-negative bounded functions adapted to $\{\mathcal{F}_n\}_{n \geq 0}$. For a martingale $(f_n)_{n \geq 0}$, its generalized fractional integral $I_\gamma f = ((I_\gamma f)_n)_{n \geq 0}$ is defined as a martingale by

$$(I_\gamma f)_n = \sum_{k=0}^n \gamma_{k-1} (f_k - f_{k-1})$$

with convention $\gamma_{-1} = \gamma_0$ and $f_{-1} = 0$.

Our definition of I_γ is based on the notion of martingale transform in the sense of Burkholder [3]. For quasi-normed spaces M_1 and M_2 of functions, we denote by $B(M_1, M_2)$ the set of all bounded martingale transforms from M_1 to M_2 , that is, $I_\gamma \in B(M_1, M_2)$ means that

$$\sup_{n \geq 0} \|(I_\gamma f)_n\|_{M_2} \leq C \sup_{n \geq 0} \|f_n\|_{M_1},$$

for all M_1 -bounded martingales $f = (f_n)_{n \geq 0}$.

Let

$$\beta_n = \sum_{B \in A(\mathcal{F}_n)} P(B) \chi_B, \quad n = 0, 1, 2, \dots \quad (1.1)$$

For $\alpha > 0$, let $\gamma_n = \beta_n^\alpha$, $n \geq 0$. Then $I_\gamma f$ is the fractional integral of f introduced in [14].

In this paper we prove $I_\gamma \in B(L_\Phi, L_\Psi)$ for the Orlicz spaces L_Φ and L_Ψ under suitable conditions. Moreover, we consider the commutator $[b, I_\gamma]$ generated by a function b . For $f \in L_\infty$, which is regarded as an L_∞ -bounded martingale $f = (f_n)_{n \geq 0}$ with $f_n = E_n f$, $((I_\gamma f)_n)_{n \geq 0}$ is also an L_∞ -bounded martingale. We denote by $I_\gamma f$ the limit function, that is, $I_\gamma f = ((I_\gamma f)_n)_{n \geq 0}$. In this case the commutator $[b, I_\gamma]f = bI_\gamma f - I_\gamma(bf)$ is well-defined for all $b \in L_\infty$. In this paper we prove that, for functions b in Campanato spaces and $f \in L_\Phi$, $[b, I_\gamma]f$ is well-defined and bounded from L_Φ to L_Ψ under suitable conditions.

The definition of the Campanato space is the following:

Definition 1.2. For $p \in [1, \infty)$ and $\psi : (0, 1] \rightarrow (0, \infty)$, let

$$\mathcal{L}_{p,\psi}^- = \{f \in L_p : \|f\|_{\mathcal{L}_{p,\psi}^-} < \infty\},$$

where

$$\|f\|_{\mathcal{L}_{p,\psi}^-} = \sup_{n \geq 0} \sup_{B \in A(\mathcal{F}_n)} \frac{1}{\psi(P(B))} \left(\frac{1}{P(B)} \int_B |f - E_{n-1}f|^p dP \right)^{1/p}.$$

We say that a function $\theta : (0, 1] \rightarrow (0, \infty)$ satisfies the doubling condition if there exists a positive constant C such that, for all $r, s \in (0, 1]$,

$$\frac{1}{C} \leq \frac{\theta(r)}{\theta(s)} \leq C, \quad \text{if } \frac{1}{2} \leq \frac{r}{s} \leq 2. \quad (1.2)$$

We say that θ is almost increasing (resp. almost decreasing) if there exists a positive constant C such that, for all $r, s \in (0, 1]$,

$$\theta(r) \leq C\theta(s) \quad (\text{resp. } \theta(s) \leq C\theta(r)), \quad \text{if } r < s. \quad (1.3)$$

The stochastic basis $\{\mathcal{F}_n\}_{n \geq 0}$ is said to be regular, if there exists a constant $R \geq 2$ such that

$$f_n \leq Rf_{n-1} \quad (1.4)$$

holds for all $n \geq 1$ and all nonnegative martingales $(f_n)_{n \geq 0}$.

It is known by [12, Theorem 2.9] that, if $\{\mathcal{F}_n\}_{n \geq 0}$ is regular and ψ is almost increasing, then

$$\|f\|_{\mathcal{L}_{1,\psi}^-} \leq \|f\|_{\mathcal{L}_{p,\psi}^-} \leq C_p \|f\|_{\mathcal{L}_{1,\psi}^-}. \quad (1.5)$$

2 Orlicz spaces

First we define a set $\bar{\Phi}$ of increasing functions $\Phi : [0, \infty] \rightarrow [0, \infty]$ and give some properties of functions in $\bar{\Phi}$.

For an increasing function $\Phi : [0, \infty] \rightarrow [0, \infty]$, let

$$a(\Phi) = \sup\{t \geq 0 : \Phi(t) = 0\}, \quad b(\Phi) = \inf\{t \geq 0 : \Phi(t) = \infty\},$$

with convention $\sup \emptyset = 0$ and $\inf \emptyset = \infty$. Then $0 \leq a(\Phi) \leq b(\Phi) \leq \infty$. Let $\bar{\Phi}$ be the set of all increasing functions $\Phi : [0, \infty] \rightarrow [0, \infty]$ such that

$$0 \leq a(\Phi) < \infty, \quad 0 < b(\Phi) \leq \infty, \quad (2.1)$$

$$\lim_{t \rightarrow +0} \Phi(t) = \Phi(0) = 0, \quad (2.2)$$

$$\Phi \text{ is left continuous on } [0, b(\Phi)), \quad (2.3)$$

$$\text{if } b(\Phi) = \infty, \text{ then } \lim_{t \rightarrow \infty} \Phi(t) = \Phi(\infty) = \infty, \quad (2.4)$$

$$\text{if } b(\Phi) < \infty, \text{ then } \lim_{t \rightarrow b(\Phi)-0} \Phi(t) = \Phi(b(\Phi)) (\leq \infty). \quad (2.5)$$

In what follows, if an increasing and left continuous function $\Phi : [0, \infty) \rightarrow [0, \infty)$ satisfies (2.2) and $\lim_{t \rightarrow \infty} \Phi(t) = \infty$, then we always regard that $\Phi(\infty) = \infty$ and that $\Phi \in \bar{\Phi}$.

Definition 2.1. A function $\Phi \in \bar{\Phi}$ is called a Young function (or sometimes also called an Orlicz function) if Φ is convex on $[0, b(\Phi))$.

By the convexity, any Young function Φ is continuous on $[0, b(\Phi))$ and strictly increasing on $[a(\Phi), b(\Phi)]$. Hence Φ is bijective from $[a(\Phi), b(\Phi)]$ to $[0, \Phi(b(\Phi))]$. Moreover, Φ is absolutely continuous on any closed subinterval in $[0, b(\Phi))$. That is, its derivative Φ' exists a.e. and

$$\Phi(t) = \int_0^t \Phi'(s) ds, \quad t \in [0, b(\Phi)). \quad (2.6)$$

For $\Phi, \Psi \in \bar{\Phi}$, we write $\Phi \approx \Psi$ if there exists a positive constant C such that

$$\Phi(C^{-1}t) \leq \Psi(t) \leq \Phi(Ct) \quad \text{for all } t \in [0, \infty].$$

Definition 2.2. (i) Let Φ_Y be the set of all Young functions.

(ii) Let $\bar{\Phi}_Y$ be the set of all $\Phi \in \bar{\Phi}$ such that $\Phi \approx \Psi$ for some $\Psi \in \Phi_Y$.

(iii) Let \mathcal{Y} be the set of all $\Phi \in \Phi_Y$ such that $a(\Phi) = 0$ and $b(\Phi) = \infty$.

For $\Phi \in \bar{\Phi}$, we recall the generalized inverse of Φ in the sense of O'Neil [18, Definition 1.2].

Definition 2.3. For $\Phi \in \bar{\Phi}$ and $u \in [0, \infty]$, let

$$\Phi^{-1}(u) = \begin{cases} \inf\{t \geq 0 : \Phi(t) > u\}, & u \in [0, \infty), \\ \infty, & u = \infty. \end{cases} \quad (2.7)$$

Let $\Phi \in \bar{\Phi}$. Then Φ^{-1} is finite, increasing and right continuous on $[0, \infty)$ and positive on $(0, \infty)$. If Φ is bijective from $[0, \infty]$ to itself, then Φ^{-1} is the usual inverse function of Φ . Moreover, we have the following proposition, which is a generalization of Property 1.3 in [18].

Proposition 2.1 ([22]). *Let $\Phi \in \bar{\Phi}$. Then*

$$\Phi(\Phi^{-1}(u)) \leq u \leq \Phi^{-1}(\Phi(u)) \quad \text{for all } u \in [0, \infty]. \quad (2.8)$$

For functions $P, Q : [0, \infty] \rightarrow [0, \infty]$, we write $P \sim Q$ if there exists a positive constant C such that

$$C^{-1}P(t) \leq Q(t) \leq CP(t) \quad \text{for all } t \in [0, \infty].$$

Then, for $\Phi, \Psi \in \bar{\Phi}$,

$$\Phi \approx \Psi \quad \Leftrightarrow \quad \Phi^{-1} \sim \Psi^{-1}. \quad (2.9)$$

For a Young function Φ , its complementary function is defined by

$$\tilde{\Phi}(t) = \begin{cases} \sup\{tu - \Phi(u) : u \in [0, \infty)\}, & t \in [0, \infty), \\ \infty, & t = \infty. \end{cases}$$

Then $\tilde{\Phi}$ is also a Young function, and $(\Phi, \tilde{\Phi})$ is called a complementary pair. For example, $\Phi(t) = t$, then

$$\tilde{\Phi}(t) = \begin{cases} 0, & t \in [0, 1], \\ \infty, & t \in (1, \infty]. \end{cases}$$

Definition 2.4. For a function $\Phi \in \bar{\Phi}_Y$, let

$$\begin{aligned} L_\Phi &= \{f \in L^0 : E[\Phi(\epsilon|f|)] < \infty \text{ for some } \epsilon > 0\}, \\ \|f\|_{L_\Phi} &= \inf \{\lambda > 0 : E[\Phi(|f|/\lambda)] \leq 1\}, \\ wL_\Phi &= \left\{ f \in L^0 : \sup_{t \in (0, \infty)} \Phi(t)P(\epsilon f, t) < \infty \text{ for some } \epsilon > 0 \right\}, \\ \|f\|_{wL_\Phi} &= \inf \left\{ \lambda > 0 : \sup_{t \in (0, \infty)} \Phi(t)P(f/\lambda, t) \leq 1 \right\}, \\ \text{where } P(f, t) &= P(\{\omega \in \Omega : |f(\omega)| > t\}). \end{aligned}$$

Remark 2.1. It is known that

$$\sup_{t \in (0, \infty)} \Phi(t)P(f, t) = \sup_{t \in (0, \infty)} t P(\Phi(|f|), t), \quad (2.10)$$

see [7, Proposition 4.2] for example.

Let $(\Phi, \tilde{\Phi})$ be a complementary pair of functions in Φ_Y . Then it is known that

$$t \leq \Phi^{-1}(t)\tilde{\Phi}^{-1}(t) \leq 2t, \quad t \in [0, \infty]. \quad (2.11)$$

It is also known that

$$E[|fg|] \leq 2\|f\|_{L_\Phi}\|g\|_{L_{\tilde{\Phi}}}. \quad (2.12)$$

Lemma 2.2. Let $\Phi \in \Phi_Y$. Then, for all $A \in \mathcal{F}$, its characteristic function χ_A is in wL_Φ and

$$\|\chi_A\|_{L_\Phi} = \|\chi_A\|_{wL_\Phi} = \frac{1}{\Phi^{-1}(1/P(A))}. \quad (2.13)$$

Definition 2.5. (i) A function $\Phi \in \bar{\Phi}$ is said to satisfy the Δ_2 -condition, denote $\Phi \in \bar{\Delta}_2$, if there exists a constant $C > 0$ such that

$$\Phi(2t) \leq C\Phi(t) \quad \text{for all } t > 0. \quad (2.14)$$

- (ii) A function $\Phi \in \bar{\Phi}$ is said to satisfy the ∇_2 -condition, denote $\Phi \in \bar{\nabla}_2$, if there exists a constant $k > 1$ such that

$$\Phi(t) \leq \frac{1}{2k} \Phi(kt) \quad \text{for all } t > 0. \quad (2.15)$$

- (iii) Let $\Delta_2 = \Phi_Y \cap \bar{\Delta}_2$ and $\nabla_2 = \Phi_Y \cap \bar{\nabla}_2$.

Remark 2.2. (i) $\Delta_2 \subset \mathcal{Y}$ and $\bar{\nabla}_2 \subset \bar{\Phi}_Y$ ([10, Lemma 1.2.3]).

- (ii) Let $\Phi \in \bar{\Phi}_Y$. Then $\Phi \in \bar{\Delta}_2$ if and only if $\Phi \approx \Psi$ for some $\Psi \in \Delta_2$, and, $\Phi \in \bar{\nabla}_2$ if and only if $\Phi \approx \Psi$ for some $\Psi \in \nabla_2$.

- (iii) Let $\Phi \in \Phi_Y$. Then $\Phi \in \Delta_2$ if and only if the set of simple functions is dense in L_Φ .

- (iv) Let $\Phi \in \Phi_Y$. Then Φ^{-1} satisfies the doubling condition by its concavity, that is,

$$\Phi^{-1}(u) \leq \Phi^{-1}(2u) \leq 2\Phi^{-1}(u) \quad \text{for all } u \in [0, \infty].$$

- (v) If $\Phi \in \bar{\nabla}_2$, then there exists $\theta \in (0, 1)$ such that $\Phi((\cdot)^\theta) \in \bar{\nabla}_2$ ([22, Lemma 4.5]).

3 Main results

We denote by \mathcal{M}_{L_Φ} the set of all L_Φ bounded martingales $f = (f_n)_{n \geq 0}$.

Theorem 3.1. *Let $\Phi, \Psi \in \bar{\Phi}_Y$. Assume that $u \mapsto \Psi^{-1}(u)/\Phi^{-1}(u)$ is almost decreasing and that there exists a positive constant C such that, for all $n \geq 0$,*

$$\sum_{k=0}^n (\gamma_{k-1} - \gamma_k) \Phi^{-1} \left(\frac{1}{\beta_k} \right) + \gamma_n \Phi^{-1} \left(\frac{1}{\beta_n} \right) \leq C \Psi^{-1} \left(\frac{1}{\beta_n} \right). \quad (3.1)$$

Then, for any positive constant C_Φ , there exists a positive constant C'_Φ such that, for all $f \in \mathcal{M}_{L_\Phi}$ with $f \not\equiv 0$,

$$\Psi \left(\frac{M(I_\gamma f)}{C'_\Phi \sup_{n \geq 0} \|f_n\|_{L_\Phi}} \right) \leq \Phi \left(\frac{Mf}{C_\Phi \sup_{n \geq 0} \|f_n\|_{L_\Phi}} \right). \quad (3.2)$$

Consequently, $I_\gamma \in B(L_\Phi, wL_\Psi)$. Moreover, if $\Phi \in \nabla_2$, then $I_\gamma \in B(L_\Phi, L_\Psi)$.

Next, for a function $\rho : (0, 1] \rightarrow (0, \infty)$ such that

$$\int_0^1 \frac{\rho(t)}{t} dt < \infty, \quad (3.3)$$

let

$$\gamma_n = \int_0^{\beta_n} \frac{\rho(t)}{t} dt, \quad \beta_n = \sum_{B \in A(\mathcal{F}_n)} P(B) \chi_B, \quad n = 0, 1, 2, \dots \quad (3.4)$$

In this case we denote I_γ by I_ρ , namely, for a martingale $f = (f_n)_{n \geq 0}$,

$$I_\rho f = ((I_\rho f)_n)_{n \geq 0}, \quad (I_\rho f)_n = \sum_{k=0}^n \left(\int_0^{\beta_{k-1}} \frac{\rho(t)}{t} dt \right) (f_k - f_{k-1}). \quad (3.5)$$

If $\rho(t) = \alpha t^\alpha$ and $\alpha > 0$, then $\int_0^{\beta_{k-1}} \frac{\rho(t)}{t} dt = (\beta_{k-1})^\alpha$ and I_ρ is the fractional integrals introduced by [14] as a generalization of I_α on dyadic martingales investigated in [5].

If $\{\mathcal{F}_n\}_{n \geq 0}$ is regular, that is, there exists $R \geq 2$ such that

$$E_n f \leq R E_{n-1} f \quad (3.6)$$

for all non-negative integrable function f , then the inequality $\beta_n \leq \beta_{n-1} \leq R\beta_n$ holds, see [14, Lemma 3.1]. Hence,

$$\begin{aligned} \sum_{k=0}^n (\gamma_{k-1} - \gamma_k) \Phi^{-1}(1/\beta_k) &= \sum_{k=1}^n \Phi^{-1}(1/\beta_k) \int_{\beta_k}^{\beta_{k-1}} \frac{\rho(t)}{t} dt \\ &\sim \sum_{k=1}^n \int_{\beta_k}^{\beta_{k-1}} \frac{\Phi^{-1}(1/t) \rho(t)}{t} dt \\ &= \int_{\beta_n}^{\beta_0} \frac{\Phi^{-1}(1/t) \rho(t)}{t} dt. \end{aligned}$$

That is, (3.1) is equivalent to

$$\int_0^{\beta_n} \frac{\rho(t)}{t} dt \Phi^{-1}(1/\beta_n) + \int_{\beta_n}^{\beta_0} \frac{\rho(t) \Phi^{-1}(1/t)}{t} dt \leq C \Psi^{-1}(1/\beta_n) \quad \text{for all } n \geq 0. \quad (3.7)$$

Corollary 3.2. *Let $\{\mathcal{F}_n\}_{n \geq 0}$ be regular, and let $\Phi, \Psi \in \bar{\Phi}_Y$. Assume that $u \mapsto \Psi^{-1}(u)/\Phi^{-1}(u)$ is almost decreasing and that there exists a positive constant A such that, for all $r \in (0, 1]$,*

$$\int_0^r \frac{\rho(t)}{t} dt \Phi^{-1}(1/r) + \int_r^1 \frac{\rho(t) \Phi^{-1}(1/t)}{t} dt \leq A \Psi^{-1}(1/r). \quad (3.8)$$

Then, for any positive constant C_Φ , there exists a positive constant C_1 such that, for all $f \in \mathcal{M}_{L_\Phi}$ with $f \not\equiv 0$,

$$\Psi \left(\frac{M(I_\rho f)}{C'_\Phi \sup_{n \geq 0} \|f_n\|_{L_\Phi}} \right) \leq \Phi \left(\frac{Mf}{C_\Phi \sup_{n \geq 0} \|f_n\|_{L_\Phi}} \right). \quad (3.9)$$

Consequently, $I_\rho \in B(L_\Phi, wL_\Psi)$. Moreover, if $\Phi \in \nabla_2$, then $I_\rho \in B(L_\Phi, L_\Psi)$.

For a sequence $\gamma = (\gamma_n)_{n \geq 0}$ of positive measurable functions, let

$$M_\gamma f = \sup_{n \geq 0} \gamma_n |E_n f|, \quad f \in L_1. \quad (3.10)$$

Theorem 3.3. *Let $\Phi, \Psi \in \bar{\Phi}_Y$. Assume that $u \mapsto \Psi^{-1}(u)/\Phi^{-1}(u)$ is almost decreasing and that there exists a positive constant A such that, for all $n \geq 0$,*

$$\gamma_n \Phi^{-1}(1/\beta_n) \leq A \Psi^{-1}(1/\beta_n). \quad (3.11)$$

Then, for any positive constant C_Φ , there exists a positive constant C'_Φ such that, for all $f \in L_\Phi$ with $f \not\equiv 0$,

$$\Psi \left(\frac{M_\gamma f}{C'_\Phi \|f\|_{L_\Phi}} \right) \leq \Phi \left(\frac{Mf}{C_\Phi \|f\|_{L_\Phi}} \right). \quad (3.12)$$

Consequently, M_γ is bounded from L_Φ to wL_Ψ . Moreover, if $\Phi \in \bar{\nabla}_2$, then M_γ is bounded from L_Φ to L_Ψ .

For the commutator $[b, I_\rho]f = bI_\rho f - I_\rho(bf)$, we have the following theorem.

Theorem 3.4. *Let $\psi : (0, 1] \rightarrow (0, \infty)$, and let $\Phi, \Psi \in \bar{\Phi}_Y$.*

- (i) *Assume that ψ is almost increasing and that there exists a positive constant A and a function $\Theta \in \bar{\nabla}_2$ such that, for all $n \geq 0$,*

$$\sum_{k=0}^n (\gamma_{k-1} - \gamma_k) \Phi^{-1} \left(\frac{1}{\beta_k} \right) + \gamma_n \Phi^{-1} \left(\frac{1}{\beta_n} \right) \leq A \Theta^{-1} \left(\frac{1}{\beta_n} \right), \quad (3.13)$$

$$\psi(\beta_n) \Theta^{-1} \left(\frac{1}{\beta_n} \right) \leq A \Psi^{-1} \left(\frac{1}{\beta_n} \right), \quad (3.14)$$

$$\psi(\beta_n) \gamma_{n-1} \Phi^{-1} \left(\frac{1}{\beta_n} \right) \leq A \Psi^{-1} \left(\frac{1}{\beta_n} \right). \quad (3.15)$$

If $\Phi, \Psi \in \bar{\Delta}_2 \cap \bar{\nabla}_2$, then there exist constants $\nu \in (1, \infty)$ and $C \in (0, \infty)$ such that, for all $b \in \mathcal{L}_{\nu, \psi}^-$ and all $f \in L_\Phi$,

$$\|[b, I_\gamma]f\|_{L_\Psi} \leq C \|b\|_{\mathcal{L}_{\nu, \psi}^-} \|f\|_{L_\Phi}. \quad (3.16)$$

Moreover, if $\{\mathcal{F}_n\}_{n \geq 0}$ be regular, then, for all $b \in \mathcal{L}_{1, \psi}^-$ and all $f \in L_\Phi$,

$$\|[b, I_\gamma]f\|_{L_\Psi} \leq C \|b\|_{\mathcal{L}_{1, \psi}^-} \|f\|_{L_\Phi}, \quad (3.17)$$

without the assumption (3.15).

- (ii) Conversely, let $\{\mathcal{F}_n\}_{n \geq 0}$ be regular and $\alpha > 0$. Assume that ψ satisfies the doubling condition and that there exists a positive constant A such that, for all $n \geq 0$,

$$\Psi^{-1} \left(\frac{1}{\beta_n} \right) \leq A \beta_n^\alpha \psi(\beta_n) \Phi^{-1} \left(\frac{1}{\beta_n} \right). \quad (3.18)$$

Assume also that

$$\|b\|_{\mathcal{L}_{1,\psi}^-(\mathcal{F}_0)} = \sup_{B \in A(\mathcal{F}_0)} \frac{1}{\psi(B)P(B)} \int_B |b| dP < \infty. \quad (3.19)$$

If $[b, I_\alpha]$ is bounded from L_Φ to L_Ψ with operator norm $\|[b, I_\alpha]\|_{L_\Phi \rightarrow L_\Psi}$, then b is in $\mathcal{L}_{1,\psi}^-$ and there exists a positive constant C , independently of b , such that

$$\|b\|_{\mathcal{L}_{1,\psi}^-} \leq C \left(\|[b, I_\alpha]\|_{L_\Phi \rightarrow L_\Psi} + \|b\|_{\mathcal{L}_{1,\psi}^-(\mathcal{F}_0)} \right).$$

For an almost increasing function $\psi : (0, 1] \rightarrow (0, \infty)$, we define the sharp maximal function M_ψ^\sharp by

$$M_\psi^\sharp f = \sup_{n \geq 0} \psi(\beta_n)^{-1} E_n |f - E_{n-1} f|, \quad f \in L_1, \quad (3.20)$$

with the convention $E_{-1} f = 0$. If $\psi \equiv 1$ we denote M_ψ^\sharp by M^\sharp , that is,

$$M^\sharp f = \sup_{n \geq 0} E_n |f - E_{n-1} f|. \quad (3.21)$$

Then we define the Triebel-Lizorkin-Orlicz space as follows.

Definition 3.1. For $\Phi \in \bar{\Phi}$ and $\psi : (0, 1] \rightarrow (0, \infty)$, let

$$F_{L_\Phi}^\psi = \{f \in L_1 : \|f\|_{F_{L_\Phi}^\psi} < \infty\},$$

where

$$\|f\|_{F_{L_\Phi}^\psi} = \|M_\psi^\sharp f\|_{L_\Phi}.$$

We can extend Theorem 3.4 to Triebel-Lizorkin-Orlicz spaces

References

- [1] R. Arai and E. Nakai, Commutators of Calderón-Zygmund and generalized fractional integral operators on generalized Morrey spaces, Rev. Mat. Complut. Published online; <https://doi.org/10.1007/s13163-017-0251-4>
- [2] R. Arai, E. Nakai, G. Sadasue, Fractional integrals and their commutators on martingale Orlicz spaces, in preparation.

- [3] D. L. Burkholder, Martingale transforms, *Ann. Math. Stat.*, 37 (1966), 1494–1504.
- [4] S. Chanillo, A note on commutators, *Indiana Univ. Math. J.* 31 (1982), No. 1, 7–16.
- [5] J.-A. Chao and H. Ombe, Commutators on Dyadic Martingales, *Proc. Japan Acad.*, 61, Ser. A (1985), 35–38.
- [6] Z. Hao and Y. Jiao, Fractional integral on martingale Hardy spaces with variable exponents. *Fract. Calc. Appl. Anal.* 18 (2015), No. 5, 1128–1145.
- [7] R. Kawasumi and E. Nakai, Pointwise multipliers on weak Orlicz spaces, preprint.
- [8] M. Kikuchi, On weighted weak type maximal inequalities for martingales. *Math. Inequal. Appl.* 6 (2003), No. 1, 163–175.
- [9] M. Kikuchi, Uniform boundedness of conditional expectation operators on a Banach function space. *Math. Inequal. Appl.* 16, No. 2 (2013), 483–499.
- [10] V. Kokilashvili and M. Krbeć, Weighted inequalities in Lorentz and Orlicz spaces, World Scientific Publishing Co., Inc., River Edge, NJ, 1991.
- [11] R. L. Long, Martingale spaces and inequalities, Peking University Press, Beijing, 1993. ISBN: 7-301-02069-4
- [12] T. Miyamoto, E. Nakai and G. Sadasue, Martingale Orlicz-Hardy spaces, *Math. Nachr.* 285 (2012), No. 5-6, 670–686.
- [13] E. Nakai, On generalized fractional integrals, *Taiwanese J. Math.* 5 (2001), no. 3, 587–602.
- [14] E. Nakai and G. Sadasue, Martingale Morrey-Campanato spaces and fractional integrals, *J. Funct. Spaces Appl.* 2012 (2012), Article ID 673929, 29 pages. DOI:10.1155/2012/673929
- [15] E. Nakai and G. Sadasue, Characterizations of boundedness for generalized fractional integrals on martingale Morrey spaces, *Math. Inequalities Appl.* 20 (2017), No. 4, 929–947. doi:10.7153/mia-2017-20-58
- [16] E. Nakai, G. Sadasue and Y. Sawano, Martingale Morrey-Hardy and Campanato-Hardy Spaces, *J. Funct. Spaces Appl.* 2013 (2013), Article ID 690258, 14 pages. DOI:10.1155/2013/690258
- [17] J. Neveu, Discrete-parameter martingales, North-Holland, Amsterdam, 1975. ISBN 0720428106
- [18] R. O’Neil, Fractional integration in Orlicz spaces. I., *Trans. Amer. Math. Soc.* 115 (1965), 300–328.

- [19] M. Paluszyński, Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss. *Indiana Univ. Math. J.* 44 (1995), No. 1, 1–17.
- [20] G. Sadasue, Fractional integrals on martingale Hardy spaces for $0 < p \leq 1$, *Mem. Osaka Kyoiku Univ. Ser. III Nat. Sci. Appl. Sci.* 60 (2011), no.1 1–7.
- [21] G. Sadasue, Martingale Besov spaces and martingale Triebel-Lizorkin spaces, to appear in *Sci. Math. Jpn.*
- [22] M. Shi, R. Arai and E. Nakai, Generalized fractional integral operators and their commutators with functions in generalized Campanato spaces on Orlicz spaces, *Taiwanese J. Math.* to appear.
<https://arxiv.org/abs/1812.09148>
- [23] C. Watari, Multipliers for Walsh Fourier series, *Tohoku Math. J.*, 16 (1964), 239–251.
- [24] F. Weisz, Martingale Hardy spaces and their applications in Fourier analysis, *Lecture Notes in Mathematics*, 1568, Springer-Verlag, Berlin, 1994.
ISBN: 3-540-57623-1

Ryutaro Arai
 Department of Mathematics
 Ibaraki University
 Mito, Ibaraki 310-8512, Japan
 16nm101x@vc.ibaraki.ac.jp
 araryu314159@gmail.com

Eiichi Nakai
 Department of Mathematics
 Ibaraki University
 Mito, Ibaraki 310-8512, Japan
 eiichi.nakai.math@vc.ibaraki.ac.jp

Gaku Sadasue
 Department of Mathematics
 Osaka Kyoiku University
 Kashiwara, Osaka 582-8582, Japan
 sadasue@cc.osaka-kyoiku.ac.jp